



Matthew Gotlin, MD
159 East 74th Street
New York, NY 10021

REHAB PROTOCOL: NON-OPERATIVE TREATMENT OF PROXIMAL HUMERUS FRACTURE

Name: _____ **Date:** _____

Diagnosis: _____ **Date of Surgery:** _____

Phase I (Protection Phase): Weeks 0-4

Precautions

- Patient to remain in sling for 4-6 weeks.
- **NO** pushing yourself from chair or bed with the injured arm.
- **NO** active motion of the shoulder, **NO** resisted motions of shoulder

ROM goals:

- Pain control with ice, and modalities
- Pendulum hangs
- Active hand, wrist and elbow exercises
- Scapular exercises: Scapular pinch and roll
- No canes/pulleys until 4-6 weeks post-injury, because these are active-assist exercises

Phase II (Range of Motion phase) Weeks 5-8: (fracture moving as a unit)

Precautions

- **NO** lifting anything heavier than soda bottle
- Patient has to face the pulley and **NO** pulleys behind the back
- **AVOID** sudden jerking motion or grabbing on to objects far out from you
- **NO** excessive passive stretching beyond the limits described below
- **NO** strengthening or resisted motions of shoulder until 8 weeks after injury (resisted internal or external rotation)

Exercises

- Discontinue sling
- Cleared for daily use of your arm for activities of daily living (ADL)

159 East 74th Street
New York, NY 10021
Tel: 212-737-3301 Fax: 212-734-0407

2 Overhill Rd, Suite 310
Scarsdale, NY 10583
Tel: 914-725-6970 Fax: 914-725-6972

1600 Stewart Ave, Suite 100
Westbury, NY 11590
Tel: 516-243-8506 Fax: 212-734-0407

1870 Richmond Road
Staten Island, NY 10306
Tel: 718-285-7118 Fax: 212-734-0407

161 Atlantic Avenue
Brooklyn, NY 11201
Tel: 212-737-3301 Fax: 212-734-0407

200 West 13th Street, 6th Floor
New York, NY 10077
Tel: 212-737-3301 Fax: 212-734-0407

9226 Kennedy Boulevard
North Bergen, NJ 07047
Tel: 212-737-3301 Fax: 212-734-0407

- Pain control with ice/heat, and modalities
- Begin AAROM and advanced to AROM as tolerated-(Use of pulleys, canes for ROM is allowed)
- Active hand, wrist and elbow exercises
- Scapular exercises: Scapular pinch and roll and shoulder shrug
- Isometrics with arm at side and light resistive therabands beginning at 8 weeks

Phase III (strengthening phase): Weeks 9-16

Precautions

- Patient should have full ROM, no substitution patterns and good scapulothoracic control
- **NO** heavy lifting (>10 pounds)
- Patient has to face the pulley and **NO** pulleys behind the back
- **AVOID** sudden jerking motion or sudden reaching out to grabbing objects
- Lateral raises and side raises with weights should be with the bent elbow and below the shoulder level

Exercises

- Advance to full ROM as tolerated with passive stretching at end ranges
- Begin scapular strengthening exercises, closed chain for deltoid, PRE's (progressive resistance exercises) for large muscle groups (pecs, lats, etc)
- Isometrics with arm at side and light resistive therabands beginning at 10 weeks
- Advance strengthening as tolerated: isometrics ‡ bands (progress with increasing intensity) ‡ light weights (1-5 lbs); per rotator cuff, deltoid, and scapular stabilizers
- Only do strengthening 3x/week to avoid rotator cuff tendonitis
- Begin eccentrically resisted motions and proprioception

Modalities

- ◇ Heat and Ice
- ◇ Ultrasound
- ◇ Iontophoresis
- ◇ Phonophoresis
- ◇ Therapists' discretion
- ◇ TENS
- ◇ Trigger point massage

Evaluation and others

- ◇ Teach home exercise program

Comments:

Frequency: 2 times per week

Duration: 12 weeks

Signature: _____

Date: _____