



Matthew Gotlin, MD
159 East 74th Street
New York, NY 10021

REHAB PROTOCOL: Hamstring Strain

Name: _____

Date: _____

Diagnosis: _____

Date of Surgery: _____

Phase 1

Goals

1. Protect healing tissue
2. Minimize atrophy and strength loss
3. Prevent motion loss

Protection

- Avoid excessive active or passive lengthening of the hamstrings
- Gradual progression of hip flexion and knee extension (initially begin knee extension with hip in extension)
- Progress hip flexion with knee in flexion
- Avoid antalgic gait pattern

Ice

2-3 times daily

Therapeutic exercise (performed daily)

1. Stationary bike
2. Sub maximal Isometric at 3 angles (100°, 45°, 20°)
3. Single leg balance
4. Balance Board
5. Soft tissue mobilization/Instrument assisted STM
6. Pulsed Ultrasound
7. Progressive hip strengthening
8. Pain free isotonic knee flexion
9. Sciatic nerve flossing
10. Ice with sensory electrical stimulation

159 East 74th Street
New York, NY 10021
Tel: 212-737-3301 Fax: 212-734-0407

2 Overhill Rd, Suite 310
Scarsdale, NY 10583
Tel: 914-725-6970 Fax: 914-725-6972

1600 Stewart Ave, Suite 100
Westbury, NY 11590
Tel: 516-243-8506 Fax: 212-734-0407

1870 Richmond Road
Staten Island, NY 10306
Tel: 718-285-7118 Fax: 212-734-0407

161 Atlantic Avenue
Brooklyn, NY 11201
Tel: 212-737-3301 Fax: 212-734-0407

200 West 13th Street, 6th Floor
New York, NY 10077
Tel: 212-737-3301 Fax: 212-734-0407

9226 Kennedy Boulevard
North Bergen, NJ 07047
Tel: 212-737-3301 Fax: 212-734-0407

Criteria for progression to next phase

1. Normal walking stride without pain
2. Pain-free isometric contraction against submaximal (50%-70%) resistance during prone knee flexion (90°) manual strength test

Phase 2

Goals

1. Regain pain-free hamstring strength, progressing through full range
2. Develop neuromuscular control of trunk and pelvis with progressive increase in movement speed preparing for functional movements

Protection

- Avoid end-range lengthening of hamstrings if painful

Ice

Post-exercise, 10-15 min

Therapeutic exercise (performed 5-7 d/wk)

1. Stationary bike
2. Treadmill at moderate to high intensity (progressive increasing intervals), pain-free speed and stride
3. Isokinetic eccentrics in non-lengthened state
4. Single-limb balance windmill touches without weight
5. Single leg stance with perturbation (eg ball toss, reaches)
6. Supine hamstring curls on swiss ball
7. STM/IASTM
8. Nordic hamstring drops
9. Shuttle jumps
10. Prone leg drops
11. Lateral and retro bandwalks
12. Sciatic nerve tensioning

Criteria for progression to next phase

1. Full strength (5/5) without pain during prone knee flexion (90°) manual strength test
2. Pain-free forward and backward jog, moderate intensity
3. Strength deficit less than 20% compared against uninjured limb
4. Pain free max eccentric in shortened state

Phase 3

Goals

1. Symptom-free (eg, pain and tightness) during all activities
2. Normal concentric and eccentric hamstring strength through full range of motion and speeds
3. Improve neuromuscular control of trunk and pelvis
4. Integrate postural control into sport-specific movements



Protection

- Train within symptom free intensity

Ice

Post-exercise, 10-15 min, as needed

Therapeutic exercise (performed 4-5 d/wk)

1. Treadmill moderate to high intensity as tolerated
2. Hamstring dynamic stretching
3. Isokinetic eccentric training at end ROM (ie hyperflexion)*
4. STM/IASTM
5. Plyometric jump training
6. 5-10 yard accelerations/decelerations
7. Single-limb balance windmill touches with weight on unstable surface
9. Sport-specific drills that incorporate postural control and progressive speed

Criteria for return to sport

1. Full strength without pain
2. Bilateral symmetry in knee flexion angle of peak isokinetic concentric knee flexion torque at 60°/s
2. Full range of motion without pain
3. Replication of sport specific movements at competition speed without symptoms

Comments:

Frequency: 2-3 times per week

Duration: 12 weeks

Signature: _____

Date: _____