

Matthew Gotlin, MD 159 East 74th Street New York, NY 10021

REHAB PROTOCOL: ACL,MCL, PCL Reconstruction

Name:	Date:	
Diagnosis:	Date of Surgery:	
EARLY PHASE (Weeks 0-4)		
☐ Weight Bearing and Range of Motion		
o 0-6 weeks: toe-touch weight bearing w/		
o ROM: A/AAROM 0-90 degrees as toler	rated	
☐ Brace Use:		
o Locked in full extension at all times oth	er than PT	
☐ Therapeutic Elements:		
o Modalities as needed		
o Patella Mob; SLR's with electric stim.;	co-contractions, prone nangs	
o Estim; Co-contractions o No abduction of hip or leg at any time	0	
o No prone hangs	e	
□ Goals:		
o a/aa/ROM: 0-90 degrees		
o Control pain/swelling		
o Quad control		
RECOVERY PHASE (Weeks 5-8)		
☐ Weight Bearing and Range of Motion:		
o Discontinue crutches at week 6		
☐ Brace Use:		
o At all times, open to AROM; discontinu	ue at week 8	
☐ Therapeutic Elements:		
o Continue above		
o Gentle hip abduction with no resistance o Wall-sits 0-45	below knee	
o Mini-squats with support 0-45		
o Pool therapy		
o Treadmill walking by 8 weeks		
□ Goals:		

159 East 74th Street New York, NY 10021 Tel: 212-737-3301 Fax: 212-734-0407

2 Overhill Rd, Suite 310 Scarsdale, NY 10583 Tel: 914-725-6970 Fax: 914-725-6972 1600 Stewart Ave, Suite 100 Westbury, NY 11590 Tel: 516-243-8506 Fax: 212-734-0407

Signature:	Date:	
Frequency:times per week	Duration:	weeks
Comments:		
o Return to competitive activities		
o Treadmill (walk 1-2 miles at 10- 12 min/mile pace)		
□ Goals:		
o Swimming o Begin plyometrics at 4 months		
o Large zig-zags		
o Gentle loops		
o Figure 8's		
o Slide boards o Begin agility drills		
☐ Therapeutic Elements:		
o Can start jogging/running at 6 months		
o If return to sport, fitting for custom brace by 5 months		
o None		
o Full Brace Use:		
☐ Weight Bearing and Range of Motion:		
REINTEGRATION PHASE (Months 3-5)		
o Walk 1-2 miles at 15 min/mile pace		
□ Goals:		
o Begin prone hangs and HSL		
o Stretching		
o Treadmill		
o Step-downs		
o Continue above with increased resistance		
☐ Therapeutic Elements:		
☐ Weight Bearing and Range of Motion: o Full		
STRENGTHEN PHASE (Weeks 8-12)		
0.1.0 0.1.00.10.10.10.10.10.10.10.10.10.10.10.		
o No effusion		
o a/aa/ROM: 0-110 degrees by 6 weeks and free by 8 we o SLR x 30	eeks	
	.1	