

# Matthew Gotlin, MD 159 East 74<sup>th</sup> Street New York, NY 10021

### **REHAB PROTOCOL**: Microfracture of the Femoral Condyle

Name:	Date:
Diagnosis:	Date of Surgery:

#### Phase I (Weeks 0-8)

- Weightbearing: Touchdown weightbearing (20-30% of body weight max) for 6-8 weeks o Brace required
- Range of Motion Continuous Passive Motion (CPM) Machine for 6-8 hours per day for 6-8 weeks
  - o Set CPM to 1 cycle per minute starting at level of flexion that is comfortable
  - o Advance 10° per day until full flexion is achieved
  - o Passive Range of Motion and stretching under guidance of PT
- Therapeutic Exercises
  - o Quadriceps/Hamstring isometrics
  - o Heel slides

### Phase II (Weeks 8-12)

- Weightbearing: Advance to full weightbearing as tolerated -- discontinue crutch use
- Range of Motion Advance to full/painless ROM
- Therapeutic Exercises
  - o Closed chain extension exercises
  - o Hamstring curls
  - o Toe raises
  - o Balance exercises
  - o Begin use of the stationary bicycle/elliptical

159 East 74th Street New York, NY 10021 Tel: 212-737-3301 Fax: 212-734-0407

2 Overhill Rd, Suite 310 Scarsdale, NY 10583 Tel: 914-725-6970 Fax: 914-725-6972 1600 Stewart Ave, Suite 100 Westbury, NY 11590 Tel: 516-243-8506 Fax: 212-734-0407

## **Phase III (Months 3-6)**

- Weightbearing: Full weightbearing
- Range of Motion Full/Painless ROM
- Therapeutic Exercises
  - o Advance closed chain strengthening exercises, proprioception activities
  - o Sport-specific rehabilitation
- Gradual return to athletic activity as tolerated including jumping/cutting/pivoting sports
- Maintenance program for strength and endurance

<b>Comments:</b>			
Frequency:	times per week	Duration:	weeks
Signature:		<b>Date:</b>	