



**Matthew Gotlin, MD**  
**159 East 74<sup>th</sup> Street**  
**New York, NY 10021**

**REHAB PROTOCOL: Arthroscopic SLAP Repair**

**Name:** \_\_\_\_\_

**Date:** \_\_\_\_\_

**Diagnosis:** \_\_\_\_\_

**Date of Surgery:** \_\_\_\_\_

**Phase I (Weeks 0-4)**

- Sling immobilization at all times except for showering and rehab under guidance of PT
- Range of Motion – AAROM → AROM as tolerated
  - Restrict motion to 140° of Forward Flexion, 40° of External Rotation and Internal Rotation to stomach
  - No Internal Rotation up the back/No External Rotation behind the head
- Therapeutic Exercise
  - Wrist/Hand Range of Motion
  - Grip Strengthening
  - Isometric Abduction, Internal/External Rotation exercises with elbow at side
  - No resisted Forward Flexion/Elbow Flexion (to avoid stressing the biceps origin)
- Heat/Ice before and after PT sessions

**Phase II (Weeks 4-6)**

- Discontinue sling immobilization
- Range of Motion – Increase Forward Flexion, Internal/External Rotation to full motion as tolerated
- Therapeutic Exercise
  - Advance isometrics from Phase I to use of a theraband within AROM limitations
  - Continue with Wrist/Hand Range of Motion and Grip Strengthening
  - Begin Prone Extensions and Scapular Stabilizing Exercises (traps/rhomboids/levator scapula)
  - Gentle joint mobilization
- Modalities per PT discretion

159 East 74th Street  
New York, NY 10021  
Tel: 212-737-3301 Fax: 212-734-0407

2 Overhill Rd, Suite 310  
Scarsdale, NY 10583  
Tel: 914-725-6970 Fax: 914-725-6972

1600 Stewart Ave, Suite 100  
Westbury, NY 11590  
Tel: 516-243-8506 Fax: 212-734-0407

1870 Richmond Road  
Staten Island, NY 10306  
Tel: 718-285-7118 Fax: 212-734-0407

161 Atlantic Avenue  
Brooklyn, NY 11201  
Tel: 212-737-3301 Fax: 212-734-0407

200 West 13<sup>th</sup> Street, 6<sup>th</sup> Floor  
New York, NY 10077  
Tel: 212-737-3301 Fax: 212-734-0407

9226 Kennedy Boulevard  
North Bergen, NJ 07047  
Tel: 212-737-3301 Fax: 212-734-0407

**Phase III (Weeks 6-12)**

- Range of Motion – Progress to full AROM without discomfort
- Therapeutic Exercise – Advance theraband exercises to light weights (1-5 lbs)
  - o 8-12 repetitions/2-3 sets for Rotator Cuff, Deltoid and Scapular Stabilizers
  - o Continue and progress with Phase II exercises
  - o Begin UE ergometer
- Modalities per PT discretion

**Phase IV (Months 3-6)**

- Range of Motion – Full without discomfort
- Therapeutic Exercise – Advance exercises in Phase III (strengthening 3x per week)
  - o Sport/Work specific rehabilitation
  - o Return to throwing at 4.5 months
  - o Return to sports at 6 months if approved
- Modalities per PT discretion

**Comments:**

**Frequency:** \_\_\_\_\_ **times per week**

**Duration:** \_\_\_\_\_ **weeks**

**Signature:** \_\_\_\_\_

**Date:** \_\_\_\_\_